



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the outer line followed on both wings by a brown band: the median shade dark, very dentate on the primaries; the discal spots small, black; on the outer margin some diffuse brown and whitish shades; the extreme margin finely black. Underneath pale fawn color irrorated with grayish scales; an indistinct outer row of points on the veins. Expanse, 39 mm.

Habitat: Castro, Parana.

Boarmia nebularia, sp. nov.

Wings pale fawn color, thickly irrorated with pale brownish scales; the lines very indistinct and represented rather by shades and heavier suffusion of scales; the outer line semilunular on the primaries, straight on the secondaries; a subterminal row of dark spots, more noticeable on the secondaries; a terminal row of black points. Underneath the wings are pale fawn color, thinly irrorated with brownish scales; a black point in the cells and a broad subterminal dark shade. Expanse, 35 mm.

Habitat: Petropolis, S. Brazil.

The sexes are quite similar.


ON THE TWO SPECIES OF EUDÆMONIA.

PLATES XI-XII.


BY WILLIAM BEUTENMULLER.

Several examples of *E. brachyura* and *E. argiphontes* and their larvæ, from Sierra Leone, Africa, are in the collection of Old World Heterocera of William Schaus, Esq., which was recently donated by him to the American Museum of Natural History.

Eudæmonia brachyura (Plate XI) is pink, with the spots yellowish. The larva is deep black on the upper side with the extreme sides broken with yellowish, and the under side wholly yellowish (possibly green in life). Anal plate with the two thorny spines, cervical shield and head testaceous. Along each side of the body are three rows of spines with branches of shorter spinules, on the anterior edge of the cervical shield are four short spines. Length 33 mm. August 21, 1895. *E. argiphontes* (Plate XII) is brown with pinkish shades and a dark transverse band across each wing. The larva is yellowish (probably green in life), mottled with black along the sides, cervical shield, lateral row of spines and those on 1, 2, 3, 11 and 12 segments black. Remaining rows of spines yellow (green) tipped with black, with the spinules also tipped with black. On each side of the segments 3 and 4 is a black band broken on the dorsum by the ground color. Thoracic feet brown. Length 37 mm., July.



Eudæmonia brachyura.



Eudæmonia argiphontes.